



UDC: 547.993.02:595.443.8-114.5.008

INVESTIGATION OF CHEMICAL SUBSTANCES OF TERRESTRIAL ARTHROPODS

O. M. Klyuchko

*Kavetsky Institute of Experimental Pathology, Oncology and Radiobiology of the NAS of Ukraine
45, Vasylykivska St, Kyiv 03022, Ukraine
e-mail: kelenaxx@ukr.net*

Klyuchko O.M. Investigation of chemical substances of terrestrial arthropods. **Studia Biologica**, 2019; 13(1); 129–144 • DOI: <https://doi.org/10.30970/sbi.1301.594>

Millions of *Arthropods* are inhabitants of the Earth and some biochemical products which they produce are valuable for human practice. The aim of this work was to summarize current information about chemical substances produced by *Arthropods* such as venoms, toxins and some other chemical compounds that may be used in practice. Today such products are obtained and purified using standard biochemical methods. In this article, the information about species of the terrestrial *Arthropods* – producers of substances acting on the nervous system and used in electrophysiological investigations is presented. Some versions of arthropods' toxins classification and the table with *Araneids* toxins and effects they occur are suggested as well. Main attention was paid to some *Araneidae* venoms and toxins that block synaptic transmission in glutamatergic synapses. A review of chemical composition and some biophysical properties of such venoms and toxins was suggested and their chemical structures namely of toxins from spiders of *Nephila* family and toxins from *A. lobata* venom were described. Some biophysical properties of *Araneidae* venoms and toxins are characterized. Other components of *Arthropods*' venoms were also observed. Contemporary applications of such chemical compounds in agriculture and in the environmental monitoring systems were considered as well. The obtained results evidence that the terrestrial *Arthropods* is a valuable source of chemical compounds for biophysical laboratories and their great potential has to attract more attention of investigators.

Keywords: Arthropods' venoms, toxins, receptor antagonists, transmembrane electric current

Millions of *Arthropods* are inhabiting the Earth and they can produce some biochemical products that are valuable for human practice is. Pharmacological modification by toxins from the nature an important approach in electrophysiology for membrane receptors study. For example, a significant progress has been achieved in acetylcholine receptors study due to their experimental investigations with bungarotoxin. The same phenomenon has happened to toxins obtained from one group of *Arthropods* – *Araneidae* spiders; they are considered as very promising for glutamate receptors study [16, 20, 29, 59]. It is well known that glutamatergic type of signal transmission in nervous system is widely spread in nature. It was found in many organisms, even in the phylogenetically distant ones. Toxins from *Araneidae* spiders are characterized by high specificity to glutamate channel-receptor complex (CRC) that has been achieved during millions years of evolution [16, 20, 29, 59]. This specificity makes them an invaluable tool for studying the molecular nature of numbers of physiological phenomena in animal nervous system (NS). Among the most important tasks of the physiology of synapses, one can mention mediator identification, finding of the co-relations between the mechanisms of membrane channel-receptor complex (CRC) functioning and its structure, complete understanding and adequate simulation of transmission mechanism in the motor neuronal junctions, etc. An important approach to solution of this problem was a search for antagonists that can selectively block specific types of synaptic transmission, in particular, with a certain type of membrane receptor. An interest to toxins of biological origin is constantly growing among neurophysiologists and biophysicists. These toxins have several advantages over snake toxins, such as low molecular weight and simple chemical structure. These qualities allow to conduct a targeted synthesis of homologous substances and investigate their effects on glutamate receptors.

Applications of different zootoxins for the neurophysiologic investigations were demonstrated in the monographs by professors P.G. Kostyuk, O.A. Krishtal, I.S. Magura, I.V. Skok [58, 63, 81]. Later on, these investigations were continued by the representatives of their scientific schools in collaboration with foreign colleagues [2, 36]. Nowadays, the results of studying some toxins from *Arthropodae* (including *Araneidae* toxins), as well as other similar phenol- and indole- derivatives were applied in agriculture [16, 20, 29, 59] and in ecological monitoring of environment [38–57], and were defended by patents [46–57]. It is necessary to emphasize that real increase in study these toxins happened in 1980th – mid 1990th. Because of the importance of studying *Arthropodae* toxins [1–19, 21–37, 60–97], a review is devoted to fundamental works of different authors who studied these toxins.

The present work is aimed at deep investigation of scientific literary sources about *Arthropods*' venoms, toxins and other their compounds important for biophysics, biochemistry, biotechnology, and related branches.

Venoms of the terrestrial *Arthropods* as a source of chemical compounds for biophysical laboratories. Many representatives of the Earth terrestrial and marine fauna contain poisons acting on animal nervous system. Among them there are microorganisms [66], fishes and marine animals [13, 73], snakes [25, 60, 81], and many predatory arthropods [19, 65, 77, 83, 84, 64, 96]. These organisms use poisons to kill their victims. Poisons' compositions and their chemical structure were selected to achieve the goal during million years of evolution: to influence the victim as effectively as possible.

A consequence of this was a high degree of specificity of toxins' structure, the composition of poisons, and their action at the molecular level [28]. This specificity makes them an invaluable tool for studying the molecular mechanisms of physiological processes in animal nervous system.

The study of terrestrial arthropod toxins has started in the 1980th [30, 31, 77, 91, 92]. For today, some mechanisms of toxins action on the NS of invertebrates and vertebrates have been studied [28, 77, 83, 84], and molecular structure of a number of toxins have been decoded [5, 23]. Besides, their synthesis in laboratory conditions has been done [26, 87]. These works became class because poisonous species are presented in the main groups of terrestrial arthropods, and approximately all of them produce toxins that influence specifically the nervous system (NS) [96].

The greatest significance as producers of biological toxins demonstrate three classes of the arthropods: polygons (*Myriapoda*), insects (*Insecta*) and Aranei (*Arachnida*) [77, 96]. Among the most important toxins produced by *Myriapoda*, is genus *Scolopendra* [96]. Among *Insecta*, the most studied toxins have some species from the family *Sphécidae*, mainly *Philantus triangulum* [8, 79, 89], and some *Braconidae* species [96]. A variety of toxins affecting NS is produced by the representatives of *Arachnidae* family – scorpions (*Scorpiones*) [64, 65] and spiders (*Aranei*) [14, 19, 28, 72, 88].

More than 20 000 species of spiders *Aranei* are known. They are all producing poisons, excluding *Uloboridae* family that lost their poison glands [96]. They are predators who either paralyze or kill their victims by poisons. The action of poison on victim's NS is determined by a spider is style of life. So, spider-hunters (*Labidognathous*) and jumping spiders (*Salticidae*) do not make the web (or it is very simple). They usually hunt on the leaves and vertical surfaces, and in order not to lose their victims they must instantly kill them. Therefore, they have very strong poisons with irreversible effects. The spiders that make the webs (families *Araneidae*, *Theridiidae*, and others) are the alternative. It is enough for them to paralyze their victims, often to "preserve" them as meal. Their poison may be weaker and many of its components are weakly reversible [19, 96].

The arthropods' venoms have a heterogeneous chemical composition. Some of them contain the components acting on different substrates. For example, they may cause pre-, and post-synaptic effects [37, 91]. Other types of arthropods' poisons are more homogeneous taking into account chemical structure of their components and their physiological action [12].

Classification of arthropod toxins. All toxic substances of spiders can be divided in three groups according to the mechanisms of their action.

1. Ion channel antagonists. Spiders' neurotoxins from *Segestriidae* family interact specifically with sodium channels of electroexcitable membranes of vertebrates and invertebrates. Toxins slow down inactivation, and then block sodium currents completely [83, 84]. Similar properties have some scorpion toxins [64, 65]. Toxins – antagonists of calcium channels were found in the venom of *Hololena*, *Plectreurus*, *Alegenopsis* spiders [10, 11, 28].
2. Toxins – formers of channels. They are obtained from spiders' poisons of some species of *Latrodectus* genus of *Theridiidae* family. The most known among them is *Karakurt* spider whose another name is "Black Widow" (*L. ferecedim guttatus*). Spiders of this family produce powerful proteases and toxins (neurotoxins).

They form channels for calcium, potassium, sodium in belayed phospholipid membranes. They are toxins with high molecular weight (4–100 thousands Da) and complex chemical structure [27, 83, 84, 90]. These toxins are used widely in laboratory studies, since they allow to simulate *in vitro* the work of ionic channels [15].

3. Toxins – chemoreceptors' antagonists. From spider venoms of *Araneidae* family specific antagonists for cholinoreceptors [91–94, 83, 84] were obtained as well as glutamate membranes receptors of neurons and muscle cells of vertebrates and invertebrates [1, 28, 30, 31, 83, 84, 91].

According to the place of their action, all toxins care divided into substances of pre- and post-synaptic action. The poisons of *Microbracons*, scorpions *Androctonus*, spiders *Steatoda*, *Asptoptichus*, *Latrodectus* belong to the toxins with the pre-synaptic effects [11, 64, 65, 96]. Toxins of the post-synaptic action include a number of substances isolated from poisons of spiders *Argiope*, *Araneus*, *Nephila* [12, 30, 31, 95]. However, with such distribution of close substances may be not correct. For example, the spider's poison from *Lityphantes paukillianus* increases a spontaneous mediator release from the presynaptic fibers by increasing their permeability for ions, mainly for calcium [91]. On can contrary, the toxins of *Plectreurus* spp. spiders cause a complete block of synaptic transmission due to the inhibition of calcium receipts in the presynaptic terminals [11].

Araneidae toxins that block a synaptic transmission in glutamatergic synapses. Classic example of a successful use of toxins for study of synaptic transmission and membrane receptors has been demonstrated on an example of the cholinergic synapses. Electrophysiological studies have shown that a neurotoxin from cobra binds specifically and irreversibly to postsynaptic membranes of the cholinergic synapses. This neurotoxin was used in early 1970's to isolate the cholinoreceptor from synapses of different origin [60, 81, 89]. Great achievements in the study of cholinoreceptors structure and its functions are due to a detection and use of another highly specific ligand – α -bungarotoxin and a number of other snake toxins [81].

A successful use of neurotoxins stimulated a study of zoothoxins affecting the synaptic transmission in other types of synapses. The toxins that block a transmission in glutamatergic synapses were of particular interest of investigators. Despite an intensive study of synapses of this type, very specific glutamate receptor antagonists have not been found during a long time, until the 1980th [62, 74, 75].

On early 1980th studies on the entomophags' venoms that block glutamatergic transmission appeared. These poisons were obtained in Japan from spiders *Nephila clavata* the Soviet Union from *Argiope lobata* spiders [1, 30–33] and in [23, 83, 86, 91]. The properties of these venoms and toxins were studied in details. Later, toxins with properties of glutamate receptors' antagonists were found in poisons of the *Araneus gemma*, *A. trifasciata* and other spiders [12, 28, 35]. On Table, a list of *Araneidae* species that produce antagonists of glutamatergic synapses are demonstrated. Yet this list can not be considered as complete one. Representatives of this family are widespread: *N. clavata* is an inhabitant of the Japanese islands, *N. maculata* live in Papua of New Guinea, eight species, from which *A. lobata* is the best known are inhabitants of ex-USSR territory (Caucasus, Central Asia), *Argiope aurantia*, *A. florida*, *A. trifasciata*, *Araneus gemma*, and some other species are from the New World. So, many researchers around the world can work with *Araneidae* toxins.

Toxins from *Araneidae* and their biological effectsТоксини з *Araneidae* та їхні біологічні ефекти

Species of spiders	Venoms, toxins	Effects of venoms, toxins (known or supposed)
Fam. Araneidae <i>Argiope trifasciata</i>	argiotoxin ₆₃₆ argiotoxin ₆₂₂ argiotoxin ₆₅₉ argiotoxin ₆₇₃	Dose-dependent blocking of open glutamate channels in locusts' muscles. Block of some closed channels or competitive antagonism was supposed. Allosteric modulation of glutamate binding with CRC is possible.
<i>Argiope lobata</i>	argiopin ₆₃₆	Blocking of open channels of CRC in muscles of locusts and larvae of meat fly. Possible blocking of closed channels or competitive antagonism. Blocking of open channels of nicotinous CRC in frog skeletal muscle.
<i>Araneus gemma</i> <i>Argiope florida</i> <i>Araneus andreusi</i> * [8]	the same as for <i>A. trifasciata</i>	the same as for <i>A. trifasciata</i>
<i>Argiope aurantia</i>	argiotoxin ₆₃₆	Dose-dependent blocking of non-NMDA receptors in the central nervous system of chicken. Dose-dependent blocking of transmission in glutamatergical nerve-muscle junction of locusts.
<i>Araneus diadematus</i> * [95]	Integral venom	Induction of glutamate-dependent ion currents in central nervous system of chicken.
<i>Neoscona arabeska</i>	Venom (after boiling)	Weakly reversible blocking, dose-dependent blocking in the glutamatergic neuromuscular synapse of locusts. Reverse blocking of non-NMDA transmission in central nervous system of chicken.
<i>Nephila clavata</i>	JSTX-3	Irreversible antagonist of glutamate receptors in different biological species.
<i>Nephila maculata</i>	NSTX	The expected effect is similar to JSTX-3.
Fam. Oxyopidae <i>Peucetia viridens</i>	Integral venom	Reversible blocking of transmission of non-NMDA type in the central nervous system of chicken.
Fam. Plectreuridae <i>Plectreury tristis</i>	PLTX	Long-term blockage of presynaptic calcium channels in invertebrates.
Fam. Alegenidae <i>Alegenopsis aperta</i>	AG1 AG2	Long-term blockage of calcium channels in nervous system. Reversible blockage of nerve and heart calcium channels. Anticonvulsant activity.
	T1 T	Paralyzing insect toxin: presynaptic action. Paralyzing insect toxin: presynaptic action.
<i>Hololena curta</i>	HO1 (m.w. ~ 5000) Hololena -toxin (m. w. ~ 16000)	Long-term blocking of transmission of non-NMDA type in the central nervous system of chicken. Long-term blockage of presynaptic calcium channels in invertebrates .
<i>Atrax infensus</i> * [7]	Integral poison	Swelling of tissues, increased secretion of histamine and 5-hydroxytryptamine in mammals.

Comment: Table is completed on the basis of material from [28]. Additional data are marked with *, with indication of the sources

Примітка: Таблиця укладена на основі матеріалу з [28]. Додаткові дані позначені *, зі зазначенням джерел

As antagonists of glutamatergic synapses, *Araneidae* toxins are of great value. Numerical individual toxins have been obtained and identified from different venoms. Some of them have a complex structure [12, 22].

They all are antagonists of glutamate receptors, but they differ in their characteristics of blocking. Reasonably, the application of these homologous structures of toxins allowed a progress in study of glutamate receptors. Chemical structure of some *Araneidae* toxins is shown on Fig. 1 [22, 23, 45–57]. For practical human activities, it is even more important that *Araneidae* toxins have a relatively small molecular weight – 400–700 Da and simple chemical structure [5, 12, 23]. Toxins of other arthropods are more complex, for example, the latrotoxin has a molecular weight more than 130 kDa, toxins of *Hololena* spp. spiders – 5–10 kDa [28]. This facilitates a study of *Araneidae* toxins and makes it possible to synthesize them both in laboratory and in industrial conditions [18, 26, 87]. Synthetic *Araneidae* toxins can be used in medicine and pharmacology as medical preparations, as well as pesticides in agriculture. A particular advantage of such pesticides is that the mechanism of pests populations regulating is close to natural [28, 62, 77].

Brief review of chemical composition and biophysical properties of Araneidae venoms and toxins. Chemical structures of toxins. In parallel with electrophysiological studies of the properties of Araneidae venoms and toxins, the investigators improved the methods of obtaining individual toxins from the integral venoms. For today, the molecular weights and chemical structure of different *Araneidae* toxins were investigated. For applied purposes, there were not enough to have small quantities of toxins obtained from natural sources. Until 1988, three most important and the most studied araneid toxins were synthesized: JSTX-3, NSTX-3, and argiopin (AR).

A. Toxins of spiders from the Nephila family. Japanese investigator Aramaki [5] obtained and purified a number of toxins from venoms *A. clavata*, *N. maculata* by HPLC on 1986. Venom *N. clavata* contains four components that possess toxic properties: JSTX-1, JSTX-2, JSTX-3, JSTX-4. The poison of *N. maculata* contains three components: NSTX-1, NSTX-2, NSTX-3. The main active agent from *N. clavata* venom was toxin JSTX-3 whose concentration in the venom was the highest [5]. The chemical formula of this toxin is given on Fig. 1. Its molecule consists of dihydrophenylacetat group coupled with asparagine and through it – with polyamine [5, 28]. Other toxic components differ between themselves by a structure of polyamines, for example, in JSTX-4 molecule, argiopine to JSTX-2 is coupled with a peptide bond [5]. In the venom of *N. maculata*, the main active agent is NSTX-3 (Fig. 1) that is very similar to JSTX-3 and 2,4-dihydrophenylacetylsparaginil.

B. Toxins of A. lobata venom. A number of investigations were devoted to isolation of the poisonous fractions from *A. lobata* venom containing toxins of various effects. The main active component of the venom is argiopin (AR) (Fig. 1), whose concentration was the highest [12]. Capability of this venom to block glutamate receptors was largely determined by the properties of argiopin [62].

The molecular weight of the obtained substance – argiopin – is 636 Da. Its chemical structure was decoded, and formula is shown on Fig. 1. It is a 2,4-dioxyphenylasparagin that is linked via polyamine with terminal arginine. In addition, from venom *A. lobata* at least 5 compounds were obtained (Fig. 1). They are similar to those in their biological activity. Like argiopin, they were isolated by the HPLC method [22, 23], but from other




Fig. 1 Chemical structure of some toxins from *Araneidae* [23]: *A* – Family of toxins from *A. lobata* – argiopines ARN. *B* – 1 – JSTX-3; 2 – NSTX-3; 3 – argiopin AR; 4 – PTX₄₃₃

Рис. 1. Хімічна структура деяких токсинів *Araneida* [23]: *A* – Сімейство токсинів з *A. lobata* – аргіопініни ARN; *B* – 1 – JSTX-3; 2 – NSTX-3; 3 – аргіопіні AR; 4 – PTX₄₃₃

fractions of venom. Argiopine was present in fraction 1; and fractions 3.7–0 were the individual compounds. Other components were identified as well.

Biophysical properties of Araneidae venoms and toxins. Because of a format of current publication, biophysical properties of *Araneidae* venoms need more detailed observation than we would like to present now. It should be mentioned that electrophysiological experiments demonstrated an ability of JSTX-3, NSTX-3, argiopine (argioxin₆₃₆) of binding the glutamate receptors of the post-synaptic membrane, blocking the processes, normally initiated in it by the glutamate [23, 69, 78, 92, 93]. JSTX-3 can be accumulated in the area of synaptic contacts [24].

It was shown that JSTX-3 irreversibly blocks electrical signals' transmission in the glutamatergic synapses, and argiopine (argioxin₆₃₆) can be removed with partial restoring of transmission characteristics [62, 69]. Argiopine (argioxin₆₃₆) is bound mainly by the activated glutamate CRCs. It is considered [28] that argiopine can bind to inactive CRC, than transfer it in the presence of the glutamate into the activated state with a subsequent blocking of the activated form. The results of most electrophysiological studies demonstrate that JSTX-3 and argiopine (argioxin₆₃₆) interact with glutamate receptor in a noncompetitive mechanism [8, 12, 35, 80]. Some numerical characteristics of blocking glutamate receptors by various toxins have been described [2, 36, 46–57].

The first attempt to isolate the glutamate receptor was done using a mixture of toxins from *A. lobata* with its further incorporation into a bilayered phospholipids membrane was done in 1983. Later two families of toxins have been isolated from *Araneidae* venoms that are the derivatives of phenolic and indoleacetic acids. The best-studied toxins are from the first family that include JSTX-3, NSTX-3, argiopine (argioxin₆₃₆). The *Araneidae* venoms contain complex of toxins whose concentrations are diverse in different species. The argiopine (argioxin₆₃₆) occurs in the venoms of the most *Araneidae* species [12, 22, 28].

Most antagonists of the glutamatergic synapses of *Araneidae* venoms are low molecular weight substances of 400–700 Da. Chemical structure of some of them was studied. Such toxins as JSTX-3, NSTX-3, argiopine and some analogs were synthesized [18, 26, 87] (Fig. 1).

Venoms and toxins of *N. clavata* and *A. lobata* were studied in O.O. Bogomoletz Institute of Physiology under O.O. Krishtal supervision. All abovementioned conclusions about the effects of venoms and toxins from *N. clavata* and *A. lobata* were verified, and new information was also obtained [2, 36, 46–57]. It was demonstrated that all studied *Araneidae* venoms and toxins (*N. clavata* integral venom and toxin from it JSTX-3, as well as toxins from *A. lobata* – argiopine, argiopinin 1, argiopinin 2), blocked the glutamate- and kainate-activated ionic currents in membranes of the pyramidal neurons of the rat hippocampus (Fig. 2). All antagonists decreased the maximum amplitudes of chemo-activated currents without affecting the kinetics of their activation and desensitization [2, 36, 46–57]. Description of some such obtained results because of their large volume is supposed to be given in future publications.

Other components of Arthropods venoms. A number of studies on the chemical composition of arthropods venoms is not enough, although the physiological effect of venoms is due not only to the toxins that are part of their composition.

During a study of some arthropods from Central Asia (spiders, tarantulas, scorpions, scolopendra), the activity of numerical enzymes that are the part of the venom has




Fig. 2. Toxins JSTX-3 (from *Nemphila clavata*) and AR (from *Argiope lobata*) influence on transmembrane electric currents. Toxin JSTX-3 blocked glutamate-activated (Glu, above) and kainat-activated (KK, middle) transmembrane ionic currents. After receiving the control response to KK, toxin JSTX-3 was applied by the background of the KK-activated current. Concentrations of Glu and KK were 1 mmol/l, JSTX-3 was 10^{-4} mol/l. $V_{\text{hold}} = -50$ mV [2, 36, 46–57]. Toxin argiopine (AR, below) blocked the open state of KK-activated ion channels. After receiving the control response to KK, the neuron was maintained in AR during 3 minutes, then on the background of AR the KK-solution was added. Concentrations: KK 1 mmol/l, AR 1.6×10^{-2} mol/l, $V_{\text{hold}} = -100$ mV. Toxin was removed from the membrane through 15 sec [2, 36, 46–57]. All records were done on different neurons [41]

Рис. 2. Токсини JSTX-3 (з *Nemphila clavata*) і AR (з *Argiope lobata*) впливають на трансмембранні електричні струми. Токсин JSTX-3 блокував глутамат-активовані (Glu, запис вгорі) та кайнат-активовані (KK, запис у центрі) трансмембранні іонні струми. Після отримання контрольної відповіді на KK, токсин JSTX-3 аплікували на тлі KK-активованих струмів. Концентрації Glu та KK становили 1 ммоль/л, JSTX-3 становила 10^{-4} моль/л. $V_{\text{підтр}} = -50$ мВ [2, 36, 46–57]. Токсин аргіопін (AR, внизу) блокував KK-активовані іонні канали у відкритому стані. Після отримання контрольної відповіді на KK нейрон витримували в AR протягом 3 хв, після того на тлі AR додавали розчин KK. Концентрації: KK 1 ммоль/л, AR 1.6×10^{-2} моль/л, $V_{\text{підтр}} = -100$ мВ. Токсин відмивали від мембрани через 15 с [2, 36, 46–57]. Усі записи зроблені на різних нейронах [41]

been recorded [3]. The activity of hyaluronidase and cholinesterase has been registered in all these arthropods. Phosphodiesterase was found in spiders of genus *Latrodectus*. High activity of kininases had been observed in a venom of karakurt, less active kininases were found in the venoms of scolopendra and tarantulas. In addition, ATPases, 5-nucleotidases, phospholipase A [3, 79] were found in poisonous arthropods.

Proteases in large quantities is a part of spiders' venoms [3, 96]. During the biting, the spider together with its toxins, introduces into victim body a digestive secret with

proteases. This type of digestion is called an “external digestion”. Due to abovementioned, the compositions of spider venoms obtained in different ways are different considerably: in the case of “milking” or, for example, in the case of homogenization of spider poison glands [8, 12, 96]. In addition, there is a large variability in quantity of different substances, both in individual samples and in neighboring species [12, 96] which can be determined by the environmental conditions of life [96]. In addition, in *Araneidae* venoms epinephrine, epinine, and dopamine were revealed in low concentrations. *Araneidae* venoms also contain 2,4-dihydroxyphenylacetic and 3,4-dihydroxyphenolacetic acids [74]. These acids are the main part of *Araneidae* toxins’.

A significant number of free amino acids, mainly glutamate [8, 12], have been registered in venoms of *Araneidae* species from New World, whose concentration can reach 130–425 mM in venom [17]. In the order of concentration decreasing in the venoms, the amino acids can be arranged as [8]:

L-glutamate → arginine → lysine → alanine → tryptophan → glycine

Thus, the antagonists of glutamatergic synapses, *Araneidae* toxins are of great value for laboratory experiments. So, *Araneidae* spiders are of interest as a source of toxins – antagonists of glutamate channel-receptor complexes. Few tens of *Araneidae* spiders species that produce different toxic substances are known nowadays [16, 20, 22, 23, 29, 59].

A number of toxins – antagonists of glutamatergic synapses have been isolated from *Araneidae* venoms. The best studied toxins among them are JSTX-3, NSTX-3, argiopine (argiotoxin₆₃₆). They are supposed to be the most potent and selective glutamate receptor antagonists [5, 62, 74]. Two families of toxins have been isolated from *Araneidae* venoms. All of them are derivatives of phenolic and indoleacetic acids. The best-studied toxins are from the first family including JSTX-3, NSTX-3, argiopine (argiotoxin₆₃₆). *Araneidae* venoms contain complex composition of toxins whose concentrations are diverse in different species. Argiopine (argiotoxin₆₃₆) occurs in the venoms of the most *Araneidae* species [12, 22, 28]. Some biochemical characteristics and biophysical effects of these venoms and toxins were characterized briefly in this publication. In 1983, the first attempt to isolate the glutamate receptor was done using a mixture of toxins from *A. lobata* with its further incorporation into a belayed phospholipids membrane.

Most antagonists of glutamatergic synapses from *Araneidae* venoms are low molecular weight substances with molecular weight of 400–700 Da and chemical structure of some of them was studied. Such toxins as JSTX-3, NSTX-3, argiopine and their analogs were synthesized [18, 26, 87]. Such properties of the glutamatergic synapse antagonists from *Araneidae* venoms, such as low molecular weight and simple chemical structure, allow them to be synthesized in industry for the purpose of new pesticides and drugs production [28]. Synthetic *Araneidae* toxins can be used in medicine and pharmacology as medical in preparations, as well as pesticides in agriculture. A particular advantage of such pesticides is that the mechanism of pests populations regulating is close to natural. Nowadays, such pesticides are enough popular in agriculture and the search of new more potent analogs is continued in different countries. These toxins were used for the biotechnological tasks [41, 42].

The toxins of *Araneidae* acted similarly on different objects. Therefore, an assumption of similarity of glutamate receptors in phylogenetically distant animals has been

suggested [28]. This review should attract attention of investigators to *Arthropodae* toxins studying and their applications these can be a source of biologically active substances for scientific laboratories and medical interest.

ACKNOWLEDGEMENTS

The author is grateful to people with whom this long-lasting research became possible: to my mother – prof. of entomology Z.F. Klyuchko (I.I. Schmalhausen Institute of Zoology of the National Academy of Sciences of Ukraine), and to my great teachers in neurophysiology – prof. O.A. Krishtal (A.A. Bogomoletz Institute of Physiology of the National Academy of Sciences of Ukraine) and prof. N. Akaike (Tokyo University, Japan), as well as to my unforgettable colleagues A.Ya. Tsyndrenko and N.I. Kiskin

The author of this study confirm that the research and publication of the results were not associated with any conflicts regarding commercial or financial relations, relations with organizations and/or individuals who may have been related to the study, and interrelations of co-authors of the article.

1. Abe T., Miwa A. Effects of a spider toxin on the glutaminergic synapse of lobster muscle. **J. Physiol.**, 1988; 389: 243–252.
2. Akaike N., Kawai N., Kiskin N.I., Klyuchko E.M., Krishtal O. A., Tsyndrenko A. Ya. Spider toxin blocks excitatory amino acid responses in isolated hippocampal pyramidal neurons. **Neurosci. Lett.**, 1987; 79: 326–330.
3. Akhunov A., Chernetsky I. I., Sadykov A.S. Biochemical characteristics of some arthropods venoms of Central Asia. **Dokl. AN USSR**, 1985; 285(4): 1009–1011. (In Russian)
4. Antonov S. M., Grishin E. V., Magazanik L. G., Shupliakov O. V., Vesselkin N. P., Volkova T.M. Argiopin blocks the glutamate responses and sensomotor transmission in motoneurons of isolated frog spinal cord. **Neurosci. Lett.**, 1987; 83(1,2): 179–184.
5. Aramaki Y., Yashuhara T., Higashijima T., Yoshioka M., Miwa A., Kawai N., Nakajima T. Chemical characterization of spider toxins JSTX and NSTX. **Proc. Japan Academy**, 1986; 62(9): 1012–1014.
6. Ashe J.H., Cox C.L., Adams M.E. Argiotoxin 636 blocks excitatory synaptic transmission in rat hippocampal neurons. **Brain Res.**, 1989; 480(1/2): 234–241.
7. Atkinson P. K. Some studies on the oedematogenic action of the venom of funnel web spiders (*Atrax species*). **Aust. J. Exp. Biol. Med. Sci.**, 1986; 64(5): 453–464.
8. Bateman A., Boden P., Dell A., Duce I.R., Quicke D.L., Usherwood P.N.R. Postsynaptic block of a glutaminergic synapse by low molecular weight fraction of spider venom. **Brain Res.**, 1985; 339(2): 237–244.
9. Boden P., Bycroft B.W., Chabra S.R., Chiplin J., Crowley P.J., Grout R.J., King T.J., McDonald E., Rafferty P., Usherwood P.N.R. The action of natural and synthetic isomers of quisqualic acid at a well-defined glutamatergic synapse. **Brain Res.**, 1986; 385(2): 205–211.
10. Bowers C., Phillips H.S., Lee P., Jan I.N., Jan L.Y. Identification and purification of an irreversible presynaptic neurotoxin from the venom of the spider *Hololena curta*. **Proc. Natl. Acad. Sci. USA**, 1987; 84(10): 8506–8510.
11. Branton W.D., Kolton L., Jan I. N., Jan L.I. Neurotoxins from *Plectreurus* spider venom are potent presynaptic blockers in *Drosophyla*. **J. Neurosci.**, 1987; 7(12): 4195–4200.
12. Budd T., Clinton P., Dell A., Duce I.R., Johnson S. J., Quicke D.L.J., Usherwood P.N.R., Ushoh G. Isolation and characterisation of glutamate receptor antagonists from venoms of orb-web spiders. **Brain Res.**, 1988; 448(2): 30–39.
13. Catterall W. Neurotoxins as allosteric modifiers of voltage-sensitive sodium channels. In: **Advances in cytopharmacology**. New York: Raven Press, 1979: 305–316.

14. Cavalieri M., D'Urso I., Lassa A., Pierdominici E., Robello H., Grusso A. Characterisation and some properties of the venom gland extract of a theriid spider (*Steatoda paukulliana*) frequently mistaken for black widow spider (*Latrodectus tredecimguttatus*). **Toxicon**, 1987; 25(9): 965–974.
15. Chanturia A.N., Mironov S.L., Sokolov Yu.V. The energy profile of the channel formed by latrotoxin in bilayer phospholipid membrane. **Ukr. Biochem. Journal.**, 1986; 58(1): 48–56. (In Russian)
16. **Chemistry and Pharmacology. The Alkaloids.** (Ed.) G.A. Cordell, A. Brossi. USA: Academic Press, 1994: 280 p.
17. Early S.L., Michaelis E.K. Presence of protein and glutamate as major constituents of the venom of the spider *Araneus gemma*. **Toxicon**, 1987; 25(4): 433–442.
18. Elin E.A., Macedo V.F., Onoprienko V.V., Osokina N.E., Tikhomirova O.V. Synthesis of argiopin. **Bioogr. Chemistry**, 1988; 14(5): 704–706. (In Russian)
19. Friede T., Nentwig W. Immobilising and lethal effects of spider venoms on the cockroach and the common meal beetle. **Toxicon**, 1989; 27(3): 305–316.
20. **Fortschritte der Chemie organischer Naturstoffe.** In: **Progress in the Chemistry of Organic Natural Products.** (Ed.) W. Herz, G.W. Kirby, R.E. Moore, W. Steglich, Ch. Tamm. USA: Springer Science & Business Media, 2012; 66: 332 p.
21. Gratton K.A.F., Clark R.B., Usherwood P.N.R. Three types of glutamate receptor on junctional membrane of locust muscle fibers. **Brain Res.**, 1979; 171(2): 360–364.
22. Grishin E.V., Volkova T.M., Arseniev A.S. Antagonists of glutamate receptors from the venom of *Argiope lobata* spider. **Bioorganicheskaya chimia**, 1988; 14(7): 883–892. (In Russian).
23. Grishin E.V., Volkova T.M., Arsenyev A.S., Reshetova O.S., Onoprienko V.V., Magazanik L.G., Antonov S.M., Fedorova I.M. Structural and functional characteristics of argiopin – ion channel blocker from venom of spider *Argiope lobata*. **Bioorganicheskaya chimia**, 1986; 12(8): 1121–1124 (In Russian)
24. Hagiwara K., Aramaki Y., Shimazaki K., Kawai N., Nakajima T. Iodinated Joro toxin (JSTX-3). Its structure and binding to the lobster neuromuscular synapse. **Chem. Pharm. Bull.**, 1988; 36(3): 1233–1236.
25. Haliwell J.V., Othman J.B., Pelchen-Mattheus A., Dolly J. O. Central action of dendrotoxin: selective reduction of transient K conductance in hippocampus and binding to localized acceptors. **Proc. Natl. Acad. Sci. USA.**, 1986; 83: 493–497.
26. Hashimoto Y., Endo Y., Shudo K., Aramaki Y., Kawai N., Nakajima T. Synthesis of spider toxin JSTX-3 and its analogs. **Tetrah. Lett.**, 1987; 28(30): 3511–3514.
27. Himmelreih N.G., Pivneva T.A., Lishko V.K., Ivanov A.P. On the calcium permeability of latrotoxin-induced synaptosomes. **Ukr. Biochem. Journal**, 1987; 59(2): 39–44. (In Russian)
28. Jackson H., Usherwood F. N. R. Spider toxins as tools for dissecting elements of excitatory amino acids transmission. **Trends In Neurosci.**, 1988; 11(6): 278–283
29. Jankovic J., Albanese A., Atassi M.Z., Dolly J.O., Hallett M., Mayer N.H. **Botulinum Toxin E-Book: Therapeutic Clinical Practice and Science.** USA: Elsevier Health Sciences, 2009. 512 p.
30. Kawai N., Miwa A., Abe T. Spider venom contains specific receptor blocker of glutaminergic synapses. **Brain Res.**, 1982; 247(1): 169–171.
31. Kawai N., Miwa A., Abe T. Effect of spider toxin on glutaminergic synapses in the mammalian brain. **Biomed. Res.**, 1982; 3(3): 353–355.
32. Kawai N., Miwa A., Abe T. Block of glutamate receptors by a spider toxin. In: Maridel P., DeFendis F.V. (Ed.) **From Molecular Pharmacology to Behavior.** New York: Raven Press, 1983: 30–34.
33. Kawai N., Yamagishi S., Saito M., Furuya K. Blockade of synaptic transmission in the squid giant synapse by a spider toxin (JSTX). **Brain Res.**, 1983; 278(2): 346–349.
34. Kawai N., Miwa A., Saito M., Pan-How H., Yosioka M. Spider toxin (JSTX) action on the glutamate synapse. **J. Physiol.**, 1984; 79(4): 228–231.

35. Kerry C.J., Ramsey R.L., Sansom M.S.P., Usherwood P.N.R. Single channel studies of non-competitive antagonism of a quisqualate sensitive glutamate receptor by argitoxin₆₃₆ – a fraction isolated from orb-weaver spider venom. **Brain Res.**, 1988; 459(2): 312–327.
36. Kiskin N.I., Klyuchko E.M., Kryshchal O.A., Tsyndrenko A.Ya., Akaike N., Kawai N. Blocking action of Nephila clavata spider toxin on ionic currents activated by glutamate and its agonists in isolated hippocampal neurons. **Neurophysiology**. 1989; 21(2): 110–116. (In Russian)
37. Kits K.S., Pick T. Action of the polyamine b-philantotoxin on neuromuscular transmission in insects. **Neuropharm.**, 1986; 25(10): 1089–1093.
38. Klyuchko O.M. **Information and computer technologies in biology and medicine**. Kyiv: NAU-druk, 2008. 252 p. (In Ukrainian)
39. Klyuchko O.M. Investigation and simulation of some antagonists influence on glutamate- and kainate-activated currents in neuronal membranes of the brain. In: **Contemporary problems of biology, ecology and chemistry**. Materials of III Intl. Sci. Conference. Zaporizhzhia: Copy Art, 2012: 230 (In Ukrainian)
40. Klyuchko O.M. Monitoring of influence on organisms of some carbohydrates environmental pollutants – derivatives of phenol and indole. In: **Contemporary problems of biology, ecology and chemistry**. Materials of IV Intl. Sci. Conference. Zaporizhzhia: Copy Art, 2015: 87–89. (In Ukrainian)
41. Klyuchko O.M. Electronic expert systems for biology and medicine. **Biotechnologia acta**, 2018; 11(6): 5–28.
[DOI: <https://doi.org/10.15407/biotech11.06.005>]
42. Klyuchko O.M. Biotechnical information systems for monitoring of chemicals in environment: biophysical approach. **Biotechnologia acta**, 2019; 12(1): 5–28.
[DOI: <https://doi.org/10.15407/biotech12.01.005>]
43. Klyuchko O.M., Klyuchko Z.F. Use of technical bioinformation systems for ecological monitoring of environment pollution by toxic organic substances in ATO zone of Ukraine. In: **Zoocenosis-2015. Biodiversity and role of animals in ecosystems**. Materials of VIII Intl. Sci. Conference. Dnipro: Lira, 2015: 30–31. (In Ukrainian)
44. Klyuchko O.M., Klyuchko Z.F. Investigation of ecological effects of pollutant substances – derivatives of phenol and indole. In: **Zoocenosis-2017. Biodiversity and role of animals in ecosystems**. Materials of IX Intl. Sci. Conference. Dnipro: Lira, 2017: 12–13. (In Ukrainian)
45. Klyuchko O.M. Method of application of biotechnical monitoring system for bioindicators' accounting with biosensor and sub-system for optical registration. – Patent UA 129987 U, G01N33/00, C12Q 1/02, C12N 15/00. Priority: 27.04.2018, u201804662, Issued: 26.11.2018, **Bull.** 22, 11 p. (In Ukrainian)
46. Klyuchko O.M. Method of cells' dissociation. Patent UA 130672 U, G01N 33/00, C12Q 1/02, C12N 15/00. Priority: 27.04.18, u201804668, Issued: 26.12.2018, **Bull.** 24, 7p. (In Ukrainian)
47. Klyuchko O.M. Method of qualitative analysis of chemical substances. Patent UA 131016 U, G01N33/50, G01N21/78, C12Q 1/60. Priority: 11.05.2018, u201805174, Issued: 10.01.2019, **Bull.** 1, 9 p. (In Ukrainian)
48. Klyuchko O.M.. Method for monitoring of chemicals influence on bioorganisms in few time intervals. Patent UA 134575 U; G01N33/00, C12N 15/00, A61P 39/00. Priority: 14.12.2018, u201812443, – Issued: 27.05.2019, **Bull.** 10, 10 p. (In Ukrainian).
49. Klyuchko O.M., Biletsky A.Ya., Lizunov G.V., Shutko V.N. Method of electrical signals generating by bio-elements in technical hybrid system. Patent UA 134574 U; A01N 1/02, G01N 33/00, A61N 1/32, B82Y 30/00. Priority: 14.12.2018, u201812442, – Issued: 27.05.2019, **Bull.** 10, 10p. (In Ukrainian)
50. Klyuchko O.M., Biletsky A.Ya., Lizunov G.V., Shutko V.N. Method for application of the system of large-scale monitoring of bioobjects with possibility of their radar control. Patent UA 134576 U; МПК G01N33/00, A61B 5/05, G01N 33/50, C12Q 1/02, G01S 13/00. Priority: 14.12.2018, u201812444, – Issued: 27.05.2019, **Bull.** 10, 10 p. (In Ukrainian)
51. Klyuchko O.M., Biletsky A.Ya., Navrotskyi D.O. Method of bio-sensor test system application. Patent UA 129923 U, G01N33/00, G01N33/50, C12Q 1/02. Priority: 22.03.2018, u201802896, Issued: 26.11.2018, **Bull.** 22, 7 p. (In Ukrainian)

52. Klyuchko O.M., Biletsky A.Ya., Navrotskyi D.O. Method of application of biotechnical monitoring system with biosensor (biosensor test system). – Patent UA 132245 U; G01N33/00. Priority: 23.03.2018 u201802893, Issued: 25.02.2014, **Bull.** 4, 7 p. (In Ukrainian)
53. Klyuchko O.M., Biletsky A.Ya., Navrotskyi D.O. Method of application of biotechnical monitoring system with biosensor and sub-system for optical registration. – Patent UA 129922 U, G01N33/50. Priority: 22.03.2018, u201802894, Issued: 26.11.2018, **Bull.** 22, 10 p. (In Ukrainian)
54. Klyuchko O.M., Biletsky A.Ya., Navrotskyi D. Method of application of biotechnical monitoring system with expert subsystem and biosensor. Patent UA 131863 U; G01N33/00, C12Q 1/02, C12N 15/00. Priority: 27.04.18, u201804663, Issued: 11.02.2019, **Bull.** 3, 7p. (In Ukrainian)
55. Klyuchko O.M., Biletsky A.Ya., Navrotskyi D.A. Method of quantitative analysis of chemical substances. Patent UA 131524 U; G01N33/50, G01N21/78, C12Q 1/60, G01N33/50, G01N21/78, C12Q 1/60. Priority: 11.05.2018, u201805175, Issued: 25.01.2019, **Bull.** 2, 10 p. (In Ukrainian)
56. Klyuchko O.M., Biletsky A.Ya. Method of qualitative analysis of hydrocarbons with harmful and toxic effect on bioobjects. Patent UA 133676 U; G01N 33/50, G01N 21/78. Priority: 06.06.2018, u201806342, Issued: 25.04.2019, **Bull.** 8, 10 p. (In Ukrainian)
57. Klyuchko O.M., Biletsky A.Ya. Method of qualitative analysis of chemical substances for the influence on electrical currents in bioobjects. Patent UA134142 U; G01N 33/50, G01N 21/78, C12Q 1/60. Priority: 06.06.2019, u201806345, Issued: 10.05.2019, **Bull.** 9, 10 p. (In Ukrainian)
58. Kostyuk P.G., Kryshchal O.A. **Mechanisms of electrical excitability of the nerve cell.** M: Nauka, 1981. 208 p. (In Russian)
59. Kusano Tomonobu, Suzuki Hideyuki. **Polyamines: A Universal Molecular Nexus for Growth, Survival, and Specialized Metabolism.** USA: Springer, 2015. 336 p.
60. Lee C.Y. Recent advances in chemistry and pharmacology of snake toxins. In: **Advances in Cytopharmacology.** New York: Raven Press, 1979: 1–16.
61. Magazanik L.G., Antonov S.M., Gmiro V.E. Mechanisms of activation and blocking of the postsynaptic membrane sensitive to glutamate. **Biological Membranes**, 1984; 1(2): 130–140. (In Russian)
62. Magazanik L.G., Antonov S.M., Fedorova I.M., Grishin E.V. The action of Agriop lobata spider venom and its low molecular weight component – argiopin on postsynaptic membranes. **Biological Membranes**, 1986; 3(12): 1204–1219. (In Russian)
63. Magura I.S. **Problems of electrical excitability of neuronal membrane.** Kyiv: Naukova Dumka, 1981. 208 p. (In Russian)
64. Martin M.F., Rochad H. Large scale purification of toxins from the venom of the scorpion *Androctonus australis Hector*. **Toxicon**, 1986; 24(11–12): 1131–1139.
65. Martin M. F., Rochat H., Marchot P., Bougis P. E. Use of high performance liquid chromatography to demonstrate quantitative variation in components of venom from the scorpion *Androctonus australis Hector*. **Toxicon**, 1987; 25(5): 569–573.
66. Matsuoka I., Syuto B., Kurinara K., Kubo S. Cytotoxic action of *Clostridium botulinum* type C, toxin on neurons of central nervous system in dissociated culture. **Jap. J. Med. Sci. Biol.**, 1986; 39(5–6): 247.
67. Michaelis E.K., Michaelis M.L., Stormann T.M., Chittenden W.L., Grubbs R.D. Purification and molecular characterization of the brain synaptic membrane glutamate-binding protein. **J. Neurochem.**, 1983; 30: 1742–1747.
68. Michaelis E.K., Galton N., Early S.L. Spider venoms inhibit L-glutamate binding to brain synaptic membrane receptors. **Proc. Natl. Acad. Sci. USA.**, 1984; 81: 5571–5574.
69. Miwa A., Kawai N. Presynaptic glutamate receptor - possible involvement of K channel. **Brain Res.**, 1986; 385(1): 161–164.
70. Miwa A., Kawai N., Saito M., Pan-Hou H., Yosioka M. Effect of spider toxin (JSTX) on excitatory postsynaptic current at neuromuscular synapse of spiny lobster. **J. Neurophys.**, 1987; 58(2): 216–220.
71. Miwa A., Kawai N., Ui M. Pertussis toxin blocks presynaptic glutamate receptor – a novel “glutamate” receptor in lobster neuromuscular synapse. **Brain Res.**, 1987; 416(1): 162–165.
72. Mylecharane E.J., Spence I., Sheumack D.D., Claassens R. Howden M. E. M. Action of robustoxin, a neurotoxic polypeptide from the venom of the male funnel-web spider (*Atrax robustus*) in anaesthetized monkeys. **Toxicon**, 1989; 24(4): 481–492.

73. *Narahashi T.* Modulation of nerve membrane sodium channels by neurotoxins. In: **Advances in Cytopharmacology**. New York: Raven Press, 1979: 293–304.
74. *Pan-Hou H., Suda Y.* Molecular action mechanism of spider toxin on glutamate receptor: role of 2,4-dihydroxyphenylacetic acid in toxin molecule. **Brain Res.**, 1987; 418(1): 198–200.
75. *Pan-Hou H., Suda Y., Sumi M., Yoshioka M., Kawai N.* Inhibitory effect of 2,4-dihydroxyphenylacetyl asparagine, a common moiety of spider toxins on glutamate binding to rat brain synaptic membranes. **Neurosci. Lett.**, 1987; 81: 199–203.
76. *Pan-Hou H., Suda Y., Sumi M., Yoshioka M., Kawai N.* A spider toxin (JSTX) inhibits L-glutamate uptake by rat brain synaptosomes. **Brain Res.**, 1989; 476(2): 354–357.
77. *Piek T.* Insect venoms and toxins. In: Kerkut G. A. (Ed.) **Comprehensive Insect Physiology, Biochemistry and Pharmacology**. Oxford: Pergamon Press, 1987; 11: 595–635.
78. *Saito M, Kawai N., Miwa A., Pan-Hou H., Yoshioka M.* Spider toxin (JSTX) blocks glutamate synapse in hippocampal pyramidal neurons. **Brain Res.**, 1985; 346(2): 397–399.
79. *Schmidt J., Blum M. S., Overal W.L.* Comparative enzymology of venoms from stringing *Hymenoptera*. **Toxicon**, 1986; 24(9): 207–291.
80. *Shuplyakov O.V, Antonov S.M., Veselkin N.P., Magazanik L.G.* The ability of argiopin to block glutamatergic synapses in the frog spinal cord. **Zhurnal Evoliycion. Biohimii i Fiziologii**, 1982; 23(2): 275–276.
81. *Skock V.I., Selianko A.A., Derckach V.A.* **Neuronal cholynoreceptors**. M: Nauka, 1987. 343 p. (In Russian)
82. *Soloway S., Wilen S.H.* Improved ferric chloride test for phenols. **Anal. Chem.**, 1952; 24(6): 979–983.
83. *Tashmuhamedov B.A.* **Ion transport through biological membranes and the mechanism of physiologically active substances action**. M: Nauka, 1985. 121p. (In Russian)
84. *Tashmuchamedov B.A., Makhmudova E.M., Usmanov P.B., Kazakov I.* Reconstitution in bilayer lipid membranes of the *Crab Potamon Transcaspicum* spider venom sensitive glutamate receptors. **Gen. Physiol. Biophys.**, 1985; 4(6): 625–630.
85. *Tashmuchamedov B.A., Makhmudova E.M., Usmanov P.B., Kazakov I., Atakuziev B.U.* Isolation and reconstruction of glutamate receptors of insects on bilayer membranes. **Uzbek. Biological Journal**, 1983; 6: 57–58. (In Russian)
86. *Tashmuchamedov B.A., Usmanov P.B., Kazakov I., Kalikulov D. Yukelson L.Ya., Atakuziev B.U.* Effects of different spider venoms on artificial and biological membranes. In: Huchó F., Ovchinnicov Y. A. (Ed.) **Toxins as Tools in Neurochemistry**. Berlin: Walter de Gruyter, 1983: 311–323.
87. *Teshima T., Wakamiya T., Aramaki Y., Nakajima T., Kawai N. Shiba T.* Synthesis of a new neurotoxin NSTX-3 of Papua New Guinean spider. **Tetrah. Lett.**, 1987; 28(30): 3509–3510.
88. *Tibballs J., Sutherland S. K., Duncan A.W.* Effects of male Sydney funnel-web spider venom in a dog and cat. **Austr. Vet. J.**, 1987; 64(2): 63–64.
89. *Tuychibaev M.U., Tashmuchamedov B.A., Magazanik L.G.* New neurotoxin from the venom of the great hornet *Vespa orientalis*. **Uzbek. Biological Journal**, 1983; 6: 3–4. (In Russian)
90. *Ushkarev Yu.A., Grishin E.V.* Karakurt neurotoxin and its interaction with the rat brain receptors. **Bioorg. Chemistry**, 1986; 12(1): 71–81 (In Russian)
91. *Usmanov P.B., Kalikulov D., Shadyeva N., Tashmuchamedov B. A.* Action of *Agriope lobata* spider venom on glutamate and cholinergic synapses. **Dokl. AN USSR**, 1983; 273(4): 1017–1018. (In Russian)
92. *Usmanov P.B., Kalikulov D., Nasledov G.A., Tashmuchamedov B.A.* The effect of the venom of spider *Segestria florentina* on the mechanism of sodium channels inactivation. **Biophysics**, 1984; 30(4): 617–619. (In Russian)
93. *Usmanov P.B., Kalikulov D., Shadyeva N.G., Nenilin A.B. Tashmuchamedov B. A.* Postsynaptic blocking of glutamatergic and cholinergic synapses as a common property of *Araneidae* spider venoms. **Toxicon**, 1985; 23(3): 528–553
94. *Usmanov P. B., Tonkikh A.K., Shadyeva H., Sadykov A. A., Tashmuchamedov B. A.* Effect of *Agriope lobata* spider venom and its components on the binding of L1 H1 glutamate to locust muscle membranes. **Ukr. Biochem. Journal**, 1988; 60(3): 78 – 81. (In Russian).

95. Vycklicky L., Krusek J., Vycklicky L., Vyskochil F. Spider venom of *Araneus* opens and desensitizes glutamate channels in chicken spinal cord neurons. **Neurosci. Lett.**, 1986; 68: 227–231.
96. Zlotkin E. Toxins derived from Arthropod venoms specially affecting insects. In: Kerkut G. A., Gilbert L. I. (Ed.) **Comprehensive Insect Physiology, Biochemistry and Pharmacology**. Oxford: Pergamon Press, 1985; 10: 499–546.
97. Yoshioka M., Narai N., Pan-Hou H., Shimazaki K., Miwa A., Kawai N. Color development upon reaction of ferric ion with the toxin JSTX, a glutamate receptor blocker present in the venom gland of the spider *Nephila clavata* (Joro spider). **Toxicon**, 1988; 26(4): 414–416.

ДОСЛІДЖЕННЯ ХІМІЧНИХ РЕЧОВИН НАЗЕМНИХ ЧЛЕНИСТОНОГИХ

О. М. Ключко

*Інститут експериментальної патології,
онкології та радіобіології ім. Р. Є. Кавецького НАН України
вул. Васильківська, 45, Київ 03022, Україна
e-mail: kelenaxx@ukr.net*

Мільйони членистоногих організмів мешкають на Землі, а деякі біохімічні продукти, які вони виробляють, є цінними для практики людини. Метою цієї роботи було узагальнити поточну інформацію про речовини, які продукують членистоногі: отрути, токсини та деякі інші хімічні сполуки, які можуть бути використані в лабораторній практиці. На сьогодні такі продукти отримують і очищають за допомогою стандартних біохімічних методів. У статті представлено інформацію про види наземних членистоногих – продуцентів речовин, що діють на нервову систему та вже були застосовані в електрофізіологічних дослідженнях. Запропоновано також деякі версії класифікації токсинів членистоногих, таблицю з токсинами аранеїд і ефектами, які вони спричиняють. Основна увага приділялася деяким отрутам і токсинам *Araneidae*, які блокують синаптичну передачу в глутаматергічних синапсах, короткий огляд хімічного складу і деяких біофізичних властивостей таких отрут і токсинів, а також хімічні структури відомих токсинів із цих пулів (токсини павуків з сімейства *Nephila* і токсини з отрути *A. lobata*). Деякі біофізичні властивості отрут і токсинів *Araneidae* коротко розглянуто. Також було розглянуто інші компоненти отрут членистоногих. Описано сучасні застосування таких хімічних сполук у сільському господарстві, у методах і системах моніторингу довкілля для їхньої стандартизації. Отримані результати проведеної роботи свідчать про те, що наземні членистоногі є цінним джерелом хімічних сполук для біофізичних лабораторій, і цей великий потенціал має привертати більше уваги сучасних дослідників.

Ключові слова: отрути членистоногих, токсини, антагоністи рецепторів, трансмембранний електричний струм

Одержано: 09.05.2019